

**NON-OPIOID
EXPAREL[®]**
(bupivacaine liposome injectable suspension)

**JOIN US FOR AN IN-
PERSON PROGRAM ON
Optimizing Recovery
After Surgery with
EXPAREL**

Host: Bernadette Villamil

609-234-4025 and
bernadette.villamil@pacira.com

Thursday, November 10th, 2022

Arrival time: 6:30 PM EST

Program start time: 7:00 PM EST

RedFire Grill & Steakhouse

400 Lantana Drive
Hockessin, DE

Presented by:

Kelly Leblanc, MD

Anesthesiologist (Adult/Pediatric)
Houston, TX

The speaker has a consulting relationship with Pacira BioSciences, Inc.

The updated PhRMA Code does not allow alcohol to be served at this program.

Indication

EXPAREL[®] (bupivacaine liposome injectable suspension) is indicated for single-dose infiltration in patients aged 6 years and older to produce postsurgical local analgesia and in adults as an interscalene brachial plexus nerve block to produce postsurgical regional analgesia. Safety and efficacy have not been established in other nerve blocks.

Please see Important Safety Information on the next page.

TO REGISTER,
scan here or visit
[https://pacira.radiusdirect.net/
speaker/register.php?mid=412](https://pacira.radiusdirect.net/speaker/register.php?mid=412)



Please RSVP at least 4 business days in advance. If you have any questions about this program, please contact eNOVA at 862-294-0817 or ccarney@teamenova.com.



Important Safety Information

EXPAREL is contraindicated in obstetrical paracervical block anesthesia.

Warnings and Precautions Specific to EXPAREL

Avoid additional use of local anesthetics within 96 hours following administration of EXPAREL.

EXPAREL is not recommended for the following types or routes of administration: epidural, intrathecal, regional nerve blocks **other than interscalene brachial plexus nerve block**, or intravascular or intra-articular use.

The potential sensory and/or motor loss with EXPAREL is temporary and varies in degree and duration depending on the site of injection and dosage administered and may last for up to 5 days, as seen in clinical trials.

Warnings and Precautions for Bupivacaine-Containing Products

Central Nervous System (CNS) Reactions: There have been reports of adverse neurologic reactions with the use of local anesthetics. These include persistent anesthesia and paresthesia. CNS reactions are characterized by excitation and/or depression.

Cardiovascular System Reactions: Toxic blood concentrations depress cardiac conductivity and excitability, which may lead to dysrhythmias, sometimes leading to death.

Allergic Reactions: Allergic-type reactions (eg, anaphylaxis and angioedema) are rare and may occur as a result of hypersensitivity to the local anesthetic or to other formulation ingredients.

Chondrolysis: There have been reports of chondrolysis (mostly in the shoulder joint) following intra-articular infusion of local anesthetics, which is an unapproved use.

Methemoglobinemia: Cases of methemoglobinemia have been reported with local anesthetic use.

Full Prescribing Information is available at www.EXPAREL.com.